

Medical University of South Carolina Medical Center

Hospital Infections Disclosure Act Report

Reported by: South Carolina Department of Health and Environmental Control

Surgical Site Infection (SSI) Standardized Infection Ratio by Procedure

Data Collected: 01/01/2016 - 12/31/2016

Procedure	No. of Specific Procedures Performed ^a	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio (SIR)	95% Confidence Interval
Coronary Bypass Graft (Chest and Donor Incision)	155	1	1.37	0.73	0.036, 3.592
Coronary Bypass Graft (Chest Only Incision)	15	*	*	*	*
Abdominal Hysterectomy	257	2	2.82	0.71	0.119, 2.343
Hip Prosthesis (Replacement)	336	2	3.67	0.55	0.091, 1.799
Knee Prosthesis (Replacement)	328	4	2.00	2.00	0.636, 4.829
Colon Surgery	328	12	9.95	1.21	0.653, 2.050

a. *= Too few procedures. Reporting on too few procedures is a risk to patient confidentiality and data stability. If less than twenty surgical procedures are performed, the SIR and number of infections will be suppressed until more procedures are performed.

Medical University of South Carolina Medical Center

Central Line Associated Blood Stream Infection (CLABSI) Standardized Infection Ratio (SIR)

Data Collected: 01/01/2016 - 12/31/2016

Location ^a	No. of Central Line Days ^{b,c}	No. of Infections	No. of Predicted Infections	Standardized Infection Ratio	95% Confidence Interval
All Adult Critical Care Units	14652	14	16.531	0.847	0.482,1.387
All Adult Inpatient Wards	24119	15	22.67	0.662	0.385,1.067
All Pediatric Critical Care Units	5725	5	8.24	0.607	0.222,1.345
All Pediatric Inpatient Wards	3262	6	3.25	1.848	0.749,3.843
Adult Speciality Care	7799	9	9.25	0.972	0.474,1.785
Pediatric Specialty Care	3322	4	3.94	1.015	0.322,2.448
Neonatal Intensive Care Unit	4287	9	6.59	1.365	0.666,2.506

a. The specific patient care area in which a patient is assigned while receiving care in the healthcare facility. All adult and pediatric critical care units are combined into one SIR; all adult and pediatric inpatient wards are combined into one SIR for this report.

b. Central line days are the total number of days a central line is in place for all patients in selected hospital locations.

c. * = Too few central line days. Reporting on too few central line days is a risk to confidentiality and data stability. If there are less than fifty central line days, the SIR and number of infections will be suppressed until there are more central line days to report.

Medical University of South Carolina Medical Center

Methicillin-resistant Staphylococcus aureus bloodstream infection (MRSA BSI) LabID Event Data

Facility Wide Inpatient Data Collected: 01/01/2016 - 12/31/2016

Hospital Onset MRSA BSI Standardized Infection Ratio (SIR)				
No. Patient Days	No. LabID Events ^a	Predicted No. of LabID Events	SIR	95% Confidence Interval
215482	15	22.170863	0.677	0.393, 1.091

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)

Medical University of South Carolina Medical Center

Clostridium Difficile Infections(CDI) LabID Event Data Facility Wide Inpatient Data Collected: 01/01/2016 - 12/31/2016

Hospital Onset CDI LabID Event Data				
No. Patient Days	No. of LabID Events ^a	Predicted No. of LabID Events	SIR	95% Confidence Interval
192394	149	155.13597	0.960	0.815, 1.124

a. Hospital Onset: LabID event specimen collected as an inpatient >3 days after admission to the facility (i.e., on or after day 4)

Medical University of South Carolina Medical Center

Ventilator Associated Events(VAE) Data

Data Collected: 01/01/2016 - 12/31/2016

No. of IVAC-plus Events ^a	No. Ventilator Days	Predicted No. of IVAC-plus Events	SIR	95% Confidence Interval
85	14102	52.195702943	1.628	1.309, 2.004

a. IVAC-plus Events: All Ventilator associated events meeting the Infection-related Ventilator Associated Complications (IVAC) and Possible Ventilator-associated pneumonia (PVAP) definitions