

## South Carolina Drug Assistance Program (SC DAP) Formulary

Revised July 1, 2024

Brand Name	Generic Name	Drug Class	Additional Information
ANTIRETROVIRAL DR	UGS		
Abacavir	abacavir	NRTI	Before adding abacavir to the drug regimen, refer to the drug's full prescribing information.
Abacavir/ Lamivudine	abacavir / lamivudine	NRTI	Before adding an abacavir-containing medication to the drug regimen, refer to the drug's full prescribing information.
Abacavir / Lamivudine / Zidovudine	abacavir / lamivudine / zidovudine	NRTI	Before adding an abacavir-containing medication to the drug regimen, refer to the drug's full prescribing information.
Aptivus	tipranavir	Protease Inhibitor	
Biktarvy	bictegravir / emtricitabine / tenofovir alafenamide fumarate	Combination Treatment	Before prescribing, refer to the drug's full prescribing information.
Cabenuva	cabotegravir / rilpivirine	Combination Treatment	Additional Information for Cabenuva: Before prescribing, please review information provided at:  Additional Information for Cabenuva
Cimduo	lamivudine/tenofovir disoproxil fumarate	NRTI; NtRTI	Product carries a boxed warning for post-treatment acute exacerbations of hepatitis B.
Complera	emtricitabine / rilpivirine / tenofovir disoproxil fumarate	Combination Treatment	Indicated as a complete regimen for treatment of HIV-1 infection: 1) In patients 12 years of age and older with no antiretroviral treatment history and with HIV-1 RNA less than or equal to 100,000 copies/mL at the start of therapy and 2) In certain virologically suppressed (HIV-1 RNA <50 copies/mL) patients on a stable antiretroviral regimen at start of therapy in order to replace their current antiretroviral treatment regimen.
Delstrigo	doravirine / lamivudine/ tenofovir disoproxil fumarate	Combination Treatment	Product carries a boxed warning regarding the risk of post-treatment acute exacerbations of hepatitis B. Contraindicated when coadministered with drugs that are strong CYP3A inducers (decreases in doravirine plasma concentrations may occur which may decrease the product's effectiveness). Refer to the package insert for concurrently administered drugs to avoid. Precautions include new onset or worsening renal impairment, risk of adverse reactions or loss of virologic response due to drug interactions, bone loss and mineralization defects, and Immune Reconstitution Syndrome.
Descovy	emtricitabine / tenofovir alafenamide	NRTI; NtRTI	
Dovato	dolutegravir / lamivudine	Combination Treatment	Dovato carries a Boxed Warning for patients co-infected with hepatitis B virus (HBV) and HIV-1. Prior to initiating treatment, patients should be tested for HBV infection. The emergence of HBV variants associated with resistance to lamivudine has been reported in HIV-1-infected patients who have received lamivudine-containing antiretroviral regimens in the presence of concurrent infection with HBV. In addition, severe exacerbations of HBV have been reported in patients co-infected with HIV-1 and HBV who have discontinued lamivudine.
Edurant	rilpivirine	NNRTI	
Efavirenz	efavirenz	NNRTI	



elvitegravir / cobicistat / Genvoya entricitabine / tenofovir alafenamide Intelence etravirine NNRTI  Sentress, Isentress HD raltegravir / rilpivirine  Juluca doluetegravir / rilpivirine  Juluca doluetegravir / rilpivirine  Gombination Treatment  Juluca doluetegravir / rilpivirine  Combination Treatment  Juluca doluetegravir / rilpivirine  Combination Treatment  Safety record in pregnancy. Women with HIV who are of child age and are not interested in having children should be count by about this information. Women who become pregnant while dolutegravir should be switched to a regimen with a more we safety record in pregnancy. Women with HIV who are of child age and are not interested in having children should be count by about this information. Women who become pregnant while dolutegravir need to discuss options with their HIV providers  Kaletra lopinavir / ritonavir Protease Inhibitor  Lamivudine / Zidovudine  Rewirapine, Nevirapine  Re  Norvir ritonavir Protease Inhibitor  Combination Treatment  Indicated as a complete regimen for treatment of HIV-1 infer patients 12 years of age and older as initial therapy in those antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL, or to replace a stable antiretroviral no known substitutions associated with resistance to individ components of Odefsey.  Contraindicated when co-administered with drugs that are servitor concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently	Brand Name	Generic Name	Drug Class	Additional Information
Evotaz atazanavir / cobicistat Protease Inhibitor; Cytochrome P-450 Inhibitor Fosamprenavir Fosamprenavir Fuzeon entuvirtide Fusion Protein Inhibitor Fuzeon entuvirtide evitegravir / cobicistat / Gemvoya emtricitabine / tenofovir alafenamide Intelence etravirine NNRTI  Juluca doluetegravir / rilpivirine  Juluca doluetegravir / rilpivirine  Juluca doluetegravir / rilpivirine  Juluca doluetegravir / rilpivirine  Combination Treatment  Safety record in pregnancy. Women with HIV who are of chili age and are not interested in having children should be coun about this information. Women who become pregnant while dolutegravir need to discuss options with their HIV providers  Kaletra lopinavir / ritonavir Protease Inhibitor  Lamivudine / Jamivudine	Emtriva	emtricitabine	NRTI	
Fosamprenavir fosamprenavir Protease Inhibitor  Fuzeon enfuvirtide Fusion Protein Inhibitor Prior authorization required for an individual's first ADAP pre for this drug.  Genvoya enfuvirtide Fusion Protein Inhibitor Prior authorization required for an individual's first ADAP pre for this drug.  Genvoya entricitabine / tenofovir alafenamide Intelence etravirine NNRTI  Isentress, Isentress HD raltegravir Integrase Inhibitor  Combination Treatment affacts with this point, we HIV who wish to become pregnant and are currently receiving dolutegravir should be switched to a regimen with a more we safety record in pregnation. Women with IIV who are of this age and are not interested in having children should be count about this information. Women who become pregnant while dolutegravir need to discuss options with their HIV providers  Kaletra Iopinavir / ritonavir Protease Inhibitor  Kaletra Iopinavir / ritonavir NRTI  Lamivudine / Iamivudine / Iamivudine / Zidovudine NRTI  Norvir ritonavir Protease Inhibitor  NRTI  Odefsey emtricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment indicated as a complete regimen for treatment of HIV-1 infect patients 12 years of age and older as initial therapy in hose antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL; or to replace a stable antiretroviral to those whose are virologically suppressed (HIV-1 RNA less than per mL) for at least six months with no history of treatment in known substitutions associated with resistance to individ components of Odefsey  Prieltro doravirine NNRTI effectiveness. Refer to the package insert for concurrently.	Epivir	lamivudine	NRTI	
Fuzeon entwirtide Fusion Protein Inhibitor for this drug.  Prior authorization required for an individual's first ADAP profer this drug.  Prior authorization required for an individual's first ADAP profer this drug.  Prior authorization required for an individual's first ADAP profer this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profer for this drug.  Prior authorization required for an individual's first ADAP profers for this drug.  Prior authorization required for this drug.  Prior authorization require for this drug.  Prior authorization require for this drug.  Prior authoriz	Evotaz	atazanavir / cobicistat	•	
elvitegravir / cobicistat / entricitabine / tenofovir alafenamide  Intelence etravirine NNRTI  Juluca doluetegravir / rilpivirine Combination Treatment dolutegravir should be switched to a regimen with a more w safety record in pregnancy. Women with HIV who are of chill age and are not interested in having children should be count between the conductive of tenofovir alafenamide  Railvudine / Iamivudine /	Fosamprenavir	fosamprenavir	Protease Inhibitor	
elvitegravir / cobicistat / emtricitabine / tenofovir alafenamide   tenofovi	Fuzeon	enfuvirtide	Fusion Protein Inhibitor	Prior authorization required for an individual's first ADAP prescription
Integrase Inhibitor	Genvoya	emtricitabine / tenofovir	Combination Treatment	Tot this drag.
Juluca doluetegravir / rilpivirine	Intelence	etravirine	NNRTI	
Juluca doluetegravir / rilpivirine Combination Treatment safety record in pregnant and are currently receiving dolutegravir should be switched to a regimen with a more we safety record in pregnancy. Women with HIV who are of child age and are not interested in having children should be count about this information. Women who become pregnant while dolutegravir need to discuss options with their HIV providers.  Kaletra lopinavir / ritonavir Protease Inhibitor  Lamivudine / Zidovudine NRTI  Nevirapine, Nevirapine nevirapine nevirapine nevirapine nevirapine NRTI  Norvir ritonavir Protease Inhibitor  Odefsey emtricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment function of the patients 12 years of age and older as initial therapy in those antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL, or to replace a stable antiretroviral returnent history with HIV-1 RNA less than or 100,000 copies per mL, or to replace a stable antiretroviral now known substitutions associated with resistance to individe components of Odefsey.  Contraindicated when co-administered with drugs that are story concentrations may occur which may decrease the product's concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently effectiveness. Refer to the package insert for concurrently effectiveness.	Isentress, Isentress HD	raltegravir	Integrase Inhibitor	
Lamivudine / Zidovudine  Nevirapine, Nevirapine ER  Norvir ritonavir Protease Inhibitor  Odefsey  Combination Treatment fumarate  Protease Inhibitor  Combination Treatment function and some substitutions associated with resistance to individe components of Odefsey.  Contraindicated when co-administered with drugs that are streamly concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently	Juluca	doluetegravir / rilpivirine	Combination Treatment	Clinical information regarding dolutegravir: At this point, women with HIV who wish to become pregnant and are currently receiving dolutegravir should be switched to a regimen with a more well-defined safety record in pregnancy. Women with HIV who are of childbearing age and are not interested in having children should be counseled about this information. Women who become pregnant while on dolutegravir need to discuss options with their HIV providers.
Zidovudine  Nevirapine, Nevirapine ER  Norvir ritonavir Protease Inhibitor  Odefsey emtricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment funds who are virologically suppressed (HIV-1 RNA less than per mL) for at least six months with no history of treatment in o known substitutions associated with resistance to individe components of Odefsey.  Pifeltro  ARTI  NRTI  Indicated as a complete regimen for treatment of HIV-1 infect patients 12 years of age and older as initial therapy in those antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL; or to replace a stable antiretroviral reatment in oknown substitutions associated with resistance to individe components of Odefsey.  Contraindicated when co-administered with drugs that are so cytochrome P450 3A enzyme inducers. Decreases in doraviric concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently	Kaletra	lopinavir / ritonavir	Protease Inhibitor	
Norvir ritonavir Protease Inhibitor  Metricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment fumarate  Combination Treatment function in the components of Odefsey.  Contraindicated when co-administered with drugs that are stronger of Contraindicated when co-administered with drugs that are stronger of Contraindicated when co-administered with may decrease the product's concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently of the package in the	•	lamivudine / zidovudine	NRTI	
Odefsey  emtricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment fumarate  Combination Treatment fumarate  Combination Treatment function and the part of the patients 12 years of age and older as initial therapy in those antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL; or to replace a stable antiretroviral rethose who are virologically suppressed (HIV-1 RNA less than per mL) for at least six months with no history of treatment in no known substitutions associated with resistance to individe components of Odefsey.  Contraindicated when co-administered with drugs that are streated with the patient of the package in doraviring concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently services and older as initial therapy in those patients 12 years of age and older as initial therapy in those antiretroviral reatment in 100,000 copies per mL; or to replace a stable antiretroviral reatment history with HIV-1 RNA less than per mL) for at least six months with no history of treatment in no known substitutions associated with resistance to individe components of Odefsey.  Contraindicated when co-administered with drugs that are strongly of the package in doraviring concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently of the package insert for concurrent for the		nevirapine	NRTI	
Pifeltro  emtricitabine / rilpivirine / tenofovir alafenamide fumarate  emtricitabine / rilpivirine / tenofovir alafenamide fumarate  Combination Treatment fumarate  Combination Treatment fumarate  patients 12 years of age and older as initial therapy in those antiretroviral treatment history with HIV-1 RNA less than or 100,000 copies per mL; or to replace a stable antiretroviral rundon, one who are virologically suppressed (HIV-1 RNA less than per mL) for at least six months with no history of treatment in no known substitutions associated with resistance to individ components of Odefsey.  Contraindicated when co-administered with drugs that are so cytochrome P450 3A enzyme inducers. Decreases in doraviring concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently suppressed (HIV-1 RNA less than or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL; or to replace a stable antiretroviral rundon, or 100,000 copies per mL;	Norvir	ritonavir	Protease Inhibitor	
cytochrome P450 3A enzyme inducers. Decreases in doraviri concentrations may occur which may decrease the product's effectiveness. Refer to the package insert for concurrently	Odefsey	tenofovir alafenamide	Combination Treatment	Indicated as a complete regimen for treatment of HIV-1 infection in patients 12 years of age and older as initial therapy in those with no antiretroviral treatment history with HIV-1 RNA less than or equal to 100,000 copies per mL; or to replace a stable antiretroviral regimen in those who are virologically suppressed (HIV-1 RNA less than 50 copies per mL) for at least six months with no history of treatment failure and no known substitutions associated with resistance to individual components of Odefsey.
	Pifeltro	doravirine	NNRTI	administered drugs to avoid. Other warnings and precautions include risk of adverse reactions or loss of virologic response due to drug
Prezcobix darunavir / cobicistat Protease Inhibitor; Cytochrome P-450 Inhibitor	Prezcobix	darunavir / cobicistat		
Prezista darunavir Protease Inhibitor	Prezista	darunavir	Protease Inhibitor	



Brand Name	Generic Name	Drug Class	Additional Information
Retrovir	zidovudine	NRTI	
Reyataz	atazanavir	Protease Inhibitor	
Rukobia	fostemsavir	gp120 Attachment Inhibitor	
Selzentry	maraviroc	CCR5 Antagonist	Prior authorization required for an individual's first ADAP prescription for this drug.
Stavudine	stavudine	NRTI	
Stribild	elvitegravir / cobicistat / emtricitabine / tenofovir disoproxil fumarate	Combination Treatment	
Symfi, Symfi Lo	efavirenz / lamivudine / tenofovir disoproxil fumarate	Combination Treatment	Product carries a boxed warning for post-treatment acute exacerbations of hepatitis B.
Symtuza	darunavir / cobicistat / emtricitabine / tenofovir alafenamide	Combination Treatment	
Tivicay	dolutegravir	Integrase Inhibitor	Clinical information regarding dolutegravir: At this point, women with HIV who wish to become pregnant and are currently receiving dolutegravir should be switched to a regimen with a more well-defined safety record in pregnancy. Women with HIV who are of childbearing age and are not interested in having children should be counseled about this information. Women who become pregnant while on dolutegravir need to discuss options with their HIV providers.
Triumeq	abacavir / dolutegravir / lamivudine	Integrase Inhibitor; NRTI	Before adding an abacavir-containing medication to the drug regimen, refer to the full prescribing information. Clinical information regarding dolutegravir: At this point, women with HIV who wish to become pregnant and are currently receiving dolutegravir should be switched to a regimen with a more well-defined safety record in pregnancy. Women with HIV who are of childbearing age and are not interested in having children should be counseled about this information. Women who become pregnant while on dolutegravir need to discuss options with their HIV providers.
Trogarzo	lbalizumab-uiyk	Anti-CD4 Monoclonal Antibody; Monoclonal Antibody; Post-Attachment Inhibitor; Monoclonal Antibody	Prior authorization is required for this medication. To initiate the process, please use the link listed below to access the SC DAP Prior Authorization Request Form:  SC DAP Prior Authorization Form  Please note: Trogarzo will be dispensed from the Theratechnologies' approved network specialty pharmacy:  CVS Specialty Customer Care  800-237-2767 phone  800-323-2445 fax  customer.servicefax@cvshealth.com
Truvada	emtricitabine / tenofovir disoproxil fumarate	NRTI; NtRTI	<del></del>



Brand Name	Generic Name	Drug Class	Additional Information
Tybost	cobicistat	Cytochrome P-450 Inhibitor	Tybost should not be used with cobicistat-containing drugs such as Evotaz, Prezcobix, or Stribild. Tybost in combination with lopinavir/ritonavir or regimens containing ritonavir is not recommended due to similar effects of Tybost and ritonavir on CYP3A. Refer to the product's full prescribing information at the link below:
			Tybost Package Insert
Viracept	nelfinavir	Protease Inhibitor	
Viread	tenofovir	Antihepadnaviral, NRTI (Anti- HBV); NtRTI	
OPPORTUNISTIC AN	D CO-INFECTION DRUGS		
Acyclovir	acyclovir	Antiviral	
Amoxicillin	amoxicillin	Antibiotic	
Augmentin	amoxicillin clavulanate	Antibiotic	
Cipro	ciprofloxacin, oral	Antibiotic	
Clarithromycin	clarithromycin	Antibiotic	
Cleocin	clindamycin	Antibiotic	
Clotrimazole	clotrimazole	Antifungal	
Clotrimazole / betamethasone	clotrimazole / betamethasone topical	Antifungal	
Dapsone	dapsone	Antibiotic	
Diflucan	fluconazole	Antifungal	
Doxycycline hyclate	doxycycline hyclate	Antibiotic	
Doxycycline monohydrate	doxycycline monohydrate	Antibiotic	
Famciclovir	famciclovir	Antiviral	
Ketoconazole	ketoconazole tablets, topical	Antifungal	
Leucovorin	leucovorin	Opportunistic Infection	
Levofloxacin	levofloxacin, oral	Antibiotic	
Mepron	atovaquone	Antiprotozoal	
Metronidazole	metronidazole, oral	Antibiotic	
Moxifloxacin	moxifloxacin, oral	Antibiotic	
Nystatin	nystatin	Antifungal	
Nystatin/triamcin acetonide	nystatin / triamcinolone topical	Antifungal	
Relenza	zanamivir	Antiviral	
Ribavirin	ribavirin	Antiviral	
Sporanox	itraconazole	Antifungal	
Sulfadiazine	sulfadiazine	Antibiotic	
Sulfamethoxazole / trimethoprim	sulfamethoxazole / trimethoprim	Antibiotic	
		South Carolina Donartmont o	



DPH		(Products are	(Products are listed in alphabetical order by brand name, if the brand is available)	
Brand Name	Generic Name	Drug Class	Additional Information	
Tamiflu	oseltamivir	Antiviral		
Trimethoprim	trimethoprim	Antibiotic		
Valcyte	valganciclovir	Antiviral		
Valtrex	valacyclovir	Antiviral		
Vfend	voriconazole, oral	Antifungal		
Zithromax	azithromycin	Antibiotic		
ANTICONVULSANT	rs			
Neurontin	gabapentin	Anticonvulsant		
ANTIDEPRESSANT		A .::1		
Amitriptyline	amitriptyline	Antidepressant		
Celexa	citalopram	Antidepressant		
Cymbalta	duloxetine	Antidepressant		
Effexor XR	venlafaxine	Antidepressant		
Lexapro	escitalopram	Antidepressant		
Paxil	paroxetine	Antidepressant		
Prozac	fluoxetine	Antidepressant		
Remeron	mirtazapine	Antidepressant		
Trazodone	trazodone	Antidepressant		
Wellbutrin XL, SR	bupropion	Antidepressant		
Zoloft	sertraline	Antidepressant		
ANTIEMETIC AGEN	NTS			
Promethazine	promethazine	Antiemetic		
Ondansetron	ondansetron	Antiemetic		
ANTILIPEMIC AGE	NTS			
Crestor	rosuvastatin	Antilipemic Agent		
Pravastatin	pravastatin	Antilipemic Agent		
Zocor	simvastatin	Antilipemic Agent		
ANTITUBERCULOS	IS AGENTS			
Myambutol	ethambutol	Antitubercular Agent		
Mycobutin	rifabutin	Antitubercular Agent		



## South Carolina Drug Assistance Program (SC DAP) Formulary

Revised July 1, 2024

(Products are listed in alphabetical order by brand name, if the brand is available)

Brand Name	Generic Name	Drug Class	Additional Information
HIV-ASSOCIATED	LIPODYSTROPHY		
Egrifta SV	tesamorelin	Growth Hormone Releasing Factor	Prior authorization is required for this medication. To initiate the process, please use the link listed below to access the SC DAP Prior Authorization Request Form:  SC DAP Prior Authorization Form  Please note: Egrifta will be dispensed from the Theratechnologies' approved network specialty pharmacy:  CVS Specialty Customer Care 800-237-2767 phone
			800-323-2445 fax
			customer.servicefax@cvshealth.com
ORAL STEROIDS			
Prednisone	prednisone, oral	Steroid	
TOBACCO CESSAT South Carolina To	ION PRODUCTS bacco Quitline: 1-800-QUIT-N	NOW	
Bupropion SR	bupropion SR tablets	Tobacco Cessation	T. b
NicoDerm CQ	nicotine patch	Tobacco Cessation	Tobacco users have a better chance at quitting with a treatment regimen inclusive of medications and counseling. Visit the link below
Nicorette	nicotine polacrilex gum, lozenge	Tobacco Cessation	for information regarding tobacco cessation:
Nicotrol	nicotine inhaler, spray	Tobacco Cessation	Tobacco Cessation
Varenicline	varenicline tablets	Tobacco Cessation	

### **SC DAP DISPENSING GUIDELINES**

**Drug Coverage:** A prescription is required for any ADAP formulary drug, regardless of whether the product is a prescription drug or an over-the-counter drug.

Generic Drugs: Generic formulations of ADAP formulary drugs are dispensed when determined as most cost-effective.

Quantity: Drug quantities of one-month supplies (as authorized by the prescriber) of ADAP formulary drugs are dispensed.

#### **Formulary Acronyms:**

NNRTI: Non-Nucleoside Reverse Transcriptase Inhibitor NRTI: Nucleoside Reverse Transcriptase Inhibitor NtRTI: Nucleotide Reverse Transcriptase Inhibitor

**Prior Authorization:** The physician should complete the SC DAP Prior Authorization (PA) Form. Fax the completed PA request form and Supporting Documentation to SGRX at 313-264-0985 (fax). After approval of a client's initial prescription for a drug requiring prior authorization, it is not necessary to submit an additional prior authorization request for that drug for that individual. The PA request form is located at:

SC DAP Prior Authorization Form