



DPH Measles Outbreak Vaccination Recommendations for Children

Age	# of previous vaccine doses	MMR vaccine recommendations
0 – 6 months	0	• Vaccine is not recommended.
6 – 11 months	0	• One early dose is recommended for those in outbreak areas or traveling internationally, after discussion with their health care provider. • This dose will not count as one of the two recommended childhood MMR doses.
1 – 17 years	0	• Should receive first dose now and second dose no sooner than 28 days later.
	1	• Should receive a second dose of MMR vaccine at least 28 days after the first dose. • The second dose may be given earlier than 4 years of age, as long as both doses are given after 12 months old and are separated by at least 28 days.
	2	• Fully vaccinated; no additional doses recommended.



DPH Measles Outbreak Vaccination Recommendations for Adults

Year born	# of previous vaccine doses	MMR vaccine recommendations
Before 1957	N/A	Very likely exposed to measles as a child and considered immune; Vaccine not needed but may be given if desired.
Between 1957 – 1968	0	Should receive a first dose now followed by a second dose at least 28 days later.
	1 dose of inactivated MMR vaccine (or don't know what type they received)	
	1 dose of live attenuated (currently used type) MMR vaccine	Should receive a second dose.
After 1968	0	Should receive a first dose now followed by a second dose at least 28 days later.
	1	Should receive a second dose at least 28 days after the first dose.
	2	Fully vaccinated; no additional doses recommended.
Healthcare Personnel	Without presumptive evidence of immunity	Should get two doses at least 28 days apart.

* MMR vaccine is **not** recommended during pregnancy or for severely immunocompromised people.