

SUMMARY

Geographic Spread

South Carolina reported **sporadic activity**.

Virologic Surveillance

This week, a total of 84 influenza cases (83 positive rapid antigen detection tests; 1 lab confirmed test) were reported from 15 counties representing all four regions. The predominant circulating flu subtype and type was Influenza A(H3) and Influenza A. Since October 2, 2016, there have been 90,536 influenza cases (88,072 positive rapid antigen detection tests; 2,464 lab confirmed tests) reported.

Influenza-Like Illness Surveillance

This week, 0.14% of patient visits to sentinel providers were seen for an influenza-like illness (ILI). This is below South Carolina's baseline (3.13%). The ILI activity level was **minimal**.

Influenza-Associated Hospitalizations

There were 3 hospitalizations reported by 13 hospitals. Since October 2, 2016, there have been 2,550 influenza-associated hospitalizations reported.

Influenza-Associated Deaths

There were no lab confirmed deaths reported this week. Since October 2, 2016, there have been 94 influenza-associated deaths reported.

Contents

2 Virologic Surveillance

5 Influenza-Like Illness Surveillance

8 Hospitalizations and Deaths

10 National Influenza Surveillance

11 Influenza Surveillance Definitions

Virologic Surveillance


Lab Confirmed Influenza

Cases include positive testing via respiratory culture, RT-PCR, DFA, and IFA. Reporting is required to DHEC within 3 days via the South Carolina Infectious Disease and Outbreak Network (SCION) or DHEC 1129 card. This week, **1** lab confirmed case was reported. This compares to **2** lab confirmed cases this same week last season.

Table: Lab Confirmed Influenza Cases


Virus Type/ Subtype	No. Positive		No. Positive	
	This week	Previous week	Cumulative YTD 2016-17	Cumulative YTD 2015-16
A not subtyped	0	0	1100 (44.7%)	169 (15.2%)
A(H1N1)	0	0	64 (2.5%)	591 (53.4%)
A(H3)	1 (100%)	3 (100%)	415 (16.8%)	80 (7.3%)
B	0	0	885 (36.0%)	265 (24.0%)
Other	0	0	0	1 (0.1%)
Unknown	0	0	0	0
Total	1	3	2464	1106

Laboratory Confirmed Influenza Subtype by MMWR Week




Virologic Surveillance

Laboratory Confirmed Influenza Cases Average and Thresholds



Cumulative Influenza Case Rate/100,000 By County



Includes all lab confirmed and positive rapid cases

Virologic Surveillance


Positive Rapid Antigen Detection Tests

Providers are required to report weekly aggregate no. of positive results and influenza type to their respective regional health department. This week, **83** positive rapid tests were reported. This compares to **60** positive rapid tests reported this same week last season. The predominant influenza type was **Influenza A**.

Table: Positive Rapid Antigen Detection Tests

Virus Type	No. Positive		No. Positive	
	This week	Previous week	Cumulative YTD 2016-17	Cumulative YTD 2015-16
Influenza A	69 (83.1%)	40 (74.1%)	53553 (60.8%)	21510 (65.7%)
Influenza A/B	3 (3.6%)	0	655 (0.7%)	515 (1.6%)
Influenza B	11 (13.3%)	14 (25.9%)	33622 (38.2%)	10606 (32.3%)
Unknown	0	0	242 (0.3%)	145 (0.4%)
Total	83	54	88072	32776

Positive Rapid Influenza Tests Weekly Total By County



Highlighted areas indicate counts greater than or equal to 1

Influenza-Like Illness Surveillance


Influenza-Like Illness Surveillance Providers enrolled in the Center for Disease Control Influenza-Like Illness Network (ILINet) surveillance system report weekly aggregate no. of patient visits, and of those visits the number of patients seen for an influenza-like illness by age group (i.e. 0-4, 5-24, 25-49, 50-64, ≥ 65). This week, the ILI activity level was **minimal** and **0.14%** of patient visits to SC ILINet providers was attributed to an influenza-like illness. This is **below** the state baseline, **3.13%**. The ILI percentage for the current week last season was **1.78%**. Reports were received from providers in 7 counties, representing 3 of the 4 regions.

ILI Activity Level 2016-17 Calendar

OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
40	44	48	1	5	9	13	18	22	26	31	35
41	45	49	2	6	10	14	19	23	27	32	36
42	46	50	3	7	11	15	20	24	28	33	37
43	47	51	4	8	12	16	21	25	29	34	38
		52				17			30		39

ILI ACTIVITY LEVELS: **MINIMAL** **LOW** **MODERATE** **HIGH**

Percentage of Influenza-like Illness (ILI) Visits Reported by Sentinel Providers for Past and Current Seasons



Influenza-Like Illness Surveillance

Influenza-Like Illness Reported by Sentinel Providers

County	ILI %	County	ILI %
Abbeville	---	Greenwood	NR
Aiken	NR	Hampton	---
Allendale	---	Horry	---
Anderson	NR	Jasper	---
Bamberg	---	Kershaw	---
Barnwell	---	Lancaster	---
Beaufort	NR	Laurens	NR
Berkeley	NR	Lee	---
Calhoun	---	Lexington	---
Charleston	NR	Marion	---
Cherokee	---	Marlboro	NR
Chester	---	McCormick	---
Chesterfield	---	Newberry	---
Clarendon	---	Oconee	NR
Colleton	---	Orangeburg	---
Darlington	0.00%	Pickens	0.00%
Dillon	---	Richland	0.00%
Dorchester	NR	Saluda	0.00%
Edgefield	---	Spartanburg	NR
Fairfield	---	Sumter	0.13%
Florence	0.56%	Union	---
Georgetown	NR	Williamsburg	---
Greenville	NR	York	0.00%

NR: No reports received
 ---: No enrolled providers


ILI percentage is dependent upon the number of reporting providers and can be greatly influenced by a single provider with high numbers of ILI.

Influenza-Like Illness Surveillance

South Carolina Disease Alerting, Reporting & Tracking System

(SC-DARTS) is a collaborative network of syndromic surveillance systems within South Carolina. Currently this network contains the following data sources: SC Hospital Emergency Department (ED) chief-complaint data and Poison Control Center call data. The hospital ED syndromic surveillance system classifies ED chief complaint data into appropriate syndrome categories (i.e. Respiratory, GI, Fever, etc.). This week, the statewide percentage of ER visits with fever-flu syndrome was **5.00%**.

Percentage of Influenza-like Illness (ILI) and Fever/Flu Syndromic Visits



Influenza-Associated Hospitalizations and Deaths

Influenza-Associated Hospitalizations

are reported weekly in aggregate no. to their respective regional health departments.

This week **3** laboratory confirmed hospitalizations were reported by **13** hospitals. This compares to **4** lab confirmed hospitalizations this same week last season.


Table: Influenza Associated Hospitalizations

Age Group	No. Hospitalizations				
	This week	Previous week	Rate per 100,000*	Cumulative YTD 2016-17	Cumulative YTD 2015-16
0-4	0	0	38.7	117	190
5-17	0	0	17.1	133	128
18-49	0	0	17.3	346	448
50-64	1	2	63.2	576	476
65+	2	0	218.1	1378	621
Unknown	0	0	--	0	0
Total	3	2	55.1	2550	1866

*Population size based on 2010 Census Data

Laboratory confirmation for hospitalizations and deaths includes culture, PCR, DFA, IFA, rapid antigen detection testing, or autopsy (deaths only).

Influenza-Associated Hospitalizations and Deaths by MMWR Week



Influenza-Associated Hospitalizations and Deaths

Table: Influenza Associated Deaths

Age Group	No. Deaths				
	This week	Previous week	Rate per 100,000*	Cumulative YTD 2016-17	Cumulative YTD 2015-16
0-4	0	0	0.3	1	0
5-17	0	0	0.1	1	1
18-49	0	0	0.4	8	9
50-64	0	0	1.7	16	17
65+	0	0	10.8	68	20
Unknown	0	0	--	--	--
Total	0	0	2.0	94	47


*Population size based on 2010 Census Data

Influenza-Associated Deaths

are urgently reportable to DHEC within 24 hours of initial notification.

0 laboratory confirmed deaths were reported this week. This compares to 0 laboratory confirmed deaths this same week last season.

Influenza-Associated Deaths by Region



National Influenza Surveillance

CDC has begun summer publication of a much-abbreviated Flu View report.

To view national data, go to: <http://www.cdc.gov/flu/weekly/>

Influenza Surveillance Definitions

Activity Level (Geographic Spread): Indicator of the geographic spread of influenza activity which is reported to CDC each week.

No activity: No increase in ILI activity and no laboratory-confirmed influenza cases.

Sporadic: No increase in ILI activity and isolated laboratory-confirmed influenza cases.

Local: Increased ILI or 2 or more institutional outbreaks in one region and laboratory-confirmed influenza cases within the past 3 weeks in the region with increased ILI or outbreaks.

Regional: Increased ILI or institutional outbreaks in 2-3 regions and laboratory-confirmed influenza cases within the past 3 weeks in the regions with increased ILI or outbreaks.

Widespread: Increased ILI and/or institutional outbreaks in at least 4 regions and laboratory confirmed influenza in the state within the past 3 weeks.

Activity Level (Influenza-like Illness):

Comparison of the current week mean reported percent of visits due to ILI to the non-influenza weeks mean reported percent of visits due to ILI. The activity level corresponds to the number of standard deviations below, at, or above the mean for the current week compared to the mean of the non-influenza weeks.

Minimal: less than 2 standard deviations above the mean.

Low: 2 to less than 4 standard deviations above the mean.

Moderate: 4 to less than 6 standard deviations above the mean.

High: greater than or equal to 6 standard deviations above the mean.

Alert Threshold: indicates influenza activity (no. of laboratory confirmed influenza cases) that is higher than the past 5 seasons. Threshold is defined as 1.645 standard deviations above the average epidemic curve for each MMWR week.

Additional Resources

Center for Disease Control and Prevention
[Weekly U.S. Influenza Surveillance Report](#)

World Health Organization
[FluNet Report](#)

European Centre for Disease Prevention and Control
[Weekly Influenza Situation](#)

Government of Canada
[Weekly Influenza Report](#)

Average Epidemic Curve: Typical influenza activity (no. of lab confirmed influenza cases) for a season. Centered on the median week of the past 5 seasons, the 4 week moving average is calculated for each MMWR week.

Fever-Flu Syndrome: Includes chief complaints with any of the following ICD codes or terms: flu, fev, high temp, tem10, feel hot, night sweat, FEB, shiver, FUO, chill, 780.6, viral INF, pain all over, ILI, and body ache. Weekly fever flu count is the sum of all the records, statewide, that were categorized into the fever flu syndrome. The state denominator is a broader modification of the respiratory syndrome that includes records that have fever flu chief complaints and general respiratory illness complaints, which include: cough, coughing URI, pneumonia, croup, bronchitis, and cold. The fever flu percentage equals (weekly fever flu count/weekly state denominator)*100.

Influenza-like Illness (ILI): Fever $\geq 100^{\circ}\text{F}$ (37.8°C) AND cough AND/OR sore throat (without a known cause other than influenza). The SC baseline is the mean percentage of patient visits for ILI during non-influenza weeks (weeks when the percent of positive lab tests were less than 2% of the total season's positive lab tests for two consecutive weeks) for the previous three seasons plus two standard deviations.

MMWR week: Term for influenza surveillance week. Each week begins on Sunday and ends on Monday. Nationally, the influenza season begins with MMWR week 40 and ends with MMWR week 39. The 2016-17 influenza season began on October 2, 2016 and will end on September 30, 2017.

Seasonal Threshold: indicates the start of the influenza season when the threshold is exceeded for two consecutive weeks. Threshold is defined as the median value of the average epidemic curve.