



## This is an official **CDC Health Advisory**

Distributed via Health Alert Network  
July 17, 2021 8:45 PM  
10494-CAD-07-17-2021-MONKEYPOX

### **Potential Exposure to Person with Confirmed Human Monkeypox Infection — United States, 2021**

#### **Summary**

The Centers for Disease Control and Prevention (CDC), in collaboration with the Texas Department of State Health Services and Dallas County Health and Human Services, is investigating a single case of monkeypox virus infection in a U.S. citizen who resides in the United States and recently returned from travel to Nigeria. The patient traveled to Dallas from Lagos, Nigeria, via Atlanta on two separate flights during July 8-9, 2021. The patient presented to an emergency department in Dallas, Texas on July 13 for complaints of a rash that began on July 7, one day prior to travel. Testing at Dallas County and CDC confirmed the presence of monkeypox virus. CDC is working with the airlines to share information with state and local health officials to contact airline passengers and others who may have been in contact with the patient during two flights: Lagos, Nigeria, to Atlanta on July 8, with arrival on July 9; and Atlanta to Dallas on July 9. CDC is issuing this health advisory to ask clinicians to consider a diagnosis of monkeypox in people who present with a febrile prodrome followed by rash and who may have had direct or indirect contact with the patient.

#### **Background**

[Monkeypox](#) is endemic to several Central and West African nations. Recent cases outside of Africa either reported recent travel to one of these countries or contact with a person with confirmed monkeypox.

Symptoms of monkeypox most often begin with a prodrome of fever and other non-specific symptoms such as malaise, headache, and muscle aches following an average incubation period of 5-13 days. After the prodrome, which lasts approximately one to three days, a generalized rash appears. Nearly all patients with monkeypox have had fever early in illness onset and prior to the rash onset. Although lesions often begin on the face before spreading to other parts of the body, there has been at least one report of lesions beginning in the groin region. Lesions progress through specific stages—macules, papules, vesicles, and pustules—before scabbing and falling off<sup>1</sup>. The rash appearance of monkeypox is very similar to that of smallpox, including a centrifugal distribution and lesions on the palms and soles. Monkeypox can occur concurrently with other rash illnesses, including varicella-zoster virus and herpes simplex virus infections. Case fatality ranges between 1 and 10%. Laboratory confirmation of monkeypox is performed using real-time polymerase chain reaction (PCR) on lesion material.

A person is considered infectious beginning five days prior to rash onset and is presumed to remain infectious until lesions have crusted, those crusts have separated, and a fresh layer of skin has formed underneath. Human-to-human transmission is thought to occur primarily through large respiratory

droplets. Respiratory droplets generally cannot travel more than a few feet, so prolonged face-to-face contact is required. Transmission can also occur by direct contact with body fluids or lesion material. Indirect contact with lesion material through fomites has also been documented. Animal-to-human transmission may occur through a bite or scratch, preparation of wild game, and direct or indirect contact with body fluids or lesion material.

There is no specific treatment for monkeypox virus infection, although antivirals developed for use in patients with smallpox may prove beneficial<sup>2</sup>. Persons with direct contact (i.e., exposure to the skin, crusts, bodily fluids, or other materials) or indirect contact (e.g., presence within a 6-foot radius in the absence of an N95 or filtering respiratory for  $\geq 3$  hours) with a monkeypox patient should be monitored by health departments; some persons may be candidates for post-exposure prophylaxis with smallpox vaccine after consultation with public health authorities.

## Recommendations for Clinicians

- If clinicians identify patients with a constellation of signs and symptoms that could be monkeypox, a travel history should be solicited. Monkeypox should be considered in patients with unexplained onset of fever, chills, new rash, or new lymphadenopathy, and a history of 1) air travel from Lagos Murtala Muhammed International Airport, Nigeria, to Hartsfield-Jackson Atlanta International Airport on July 8 with arrival on July 9, 2) air travel from Atlanta to Dallas Love Field Airport on July 9, or 3) presence in those airports on July 8-9.
- Patients with suspected monkeypox should be isolated in a negative pressure room, and all personnel should wear personal protective equipment (PPE) in accordance with recommendations for standard, contact, and airborne precautions<sup>3</sup>. All healthcare workers (e.g., clinical staff and environmental staff) caring for a patient with suspect or confirmed monkeypox should be communicated the importance of maintaining proper isolation precautions so that infection is not transmitted to them or others.
- Clinicians should consult their [DHEC Regional Public Health Office](#) (see contact information below) as soon as monkeypox is suspected. DHEC will consult with CDC's monkeypox call center through the CDC Emergency Operations Center.
- After consultation with DHEC and CDC, samples can be sent to CDC or an appropriate Laboratory Response Network for confirmatory testing by PCR<sup>4</sup>. Send all specimens through the DHEC Public Health Laboratory, unless authorized to send directly to CDC.
- Ideal specimens for laboratory testing include lesion fluid, lesion roof, scabs, and crusts. Serum and whole blood can also be collected. Best practices are to collect multiple specimens from different locations on the body. Detailed specimen submission instructions are available at CDC's [monkeypox](#) website<sup>5</sup>.

## Recommendations for the Public

- Individuals who may have had contact with a suspect or confirmed monkeypox case should contact their health department for a risk assessment.

## Resources for Additional Information

1. Clinical Recognition of Monkeypox  
<https://www.cdc.gov/poxvirus/monkeypox/clinicians/clinical-recognition.html>
2. Antivirals  
<https://www.cdc.gov/poxvirus/monkeypox/clinicians/treatment.html>
3. Infection Control Measures in Hospitals  
<https://www.cdc.gov/poxvirus/monkeypox/clinicians/infection-control-hospital.html>
4. U.S. Laboratory Response Network  
<https://emergency.cdc.gov/lrn/coverage.asp>
5. Preparation and Collection of Specimens  
<https://www.cdc.gov/poxvirus/monkeypox/clinicians/prep-collection-specimens.html>

## DHEC contact information for reportable diseases and reporting requirements

Reporting of **Monkeypox** is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2021 List of Reportable Conditions available at:

<https://www.scdhec.gov/sites/default/files/Library/CR-009025.pdf>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

<b>Regional Public Health Offices – 2021</b>			
Mail or call reports to the Epidemiology Office in each Public Health Region			
<b>MAIL TO:</b>			
<b>Lowcountry</b> 4050 Bridge View Drive, Suite 600 N. Charleston, SC 29405 Fax: (843) 953-0051	<b>Midlands</b> 2000 Hampton Street Columbia, SC 29204 Fax: (803) 576-2993	<b>Pee Dee</b> 1931 Industrial Park Road Conway, SC 29526 Fax: (843) 915-6506	<b>Upstate</b> 200 University Ridge Greenville, SC 29602 Fax: (864) 282-4373
<b>CALL TO:</b>			
<b>Lowcountry</b> Allendale, Bamberg, Beaufort, Berkeley, Calhoun, Charleston, Colleton, Dorchester, Hampton, Jasper, Orangeburg  Office: (843) 441-1091 Nights/Weekends: (843) 441-1091	<b>Midlands</b> Aiken, Barnwell, Chester, Edgefield, Fairfield, Kershaw, Lancaster, Lexington, Newberry, Richland, Saluda, York  Office: (888) 801-1046 Nights/Weekends: (888) 801-1046	<b>Pee Dee</b> Clarendon, Chesterfield, Darlington, Dillon, Florence, Georgetown, Horry, Lee, Marion, Marlboro, Sumter, Williamsburg  Office: (843) 915-8886 Nights/Weekends: (843) 409-0695	<b>Upstate</b> Abbeville, Anderson, Cherokee, Greenville, Greenwood, Laurens, McCormick, Oconee, Pickens, Spartanburg, Union  Office: (864) 372-3133 Nights/Weekends: (864) 423-6648
<b>For information on reportable conditions, see</b> <a href="https://www.scdhec.gov/ReportableConditions">https://www.scdhec.gov/ReportableConditions</a>		<b>DHEC Bureau of Communicable Disease Prevention &amp; Control</b> <b>Division of Acute Disease Epidemiology</b> 2100 Bull St • Columbia, SC 29201 Phone: (803) 898-0861 • Fax: (803) 898-0897 Nights / Weekends: 1-888-847-0902	

Categories of Health Alert messages:

<b>Health Alert</b>	Conveys the highest level of importance; warrants immediate action or attention.
<b>Health Advisory</b>	Provides important information for a specific incident or situation; may not require immediate action.
<b>Health Update</b>	Provides updated information regarding an incident or situation; unlikely to require immediate action.
<b>Info Service</b>	Provides general information that is not necessarily considered to be of an emergent nature.